

## OR-7

**DIPYRAZOLODIOXADIAZOCINES AS EASILY STORABLE  
AND “READY TO USE” PRECURSORS FOR IN SITU GENERATION  
OF ENOLATE-IMINIUM 1,4-DIPOLES: AN ATOM-ECONOMICAL APPROACH  
TOPYRAZOLO[5,1-d][1,3,5]DIOXAZINES**


**V. E. Zhulanov,<sup>1</sup> P. S. Silaichev,<sup>1,2</sup> V. A. Vigovskaya,<sup>1</sup> M. V. Dmitriev<sup>1</sup>,  
A. N. Maslivets<sup>1</sup>**

<sup>1</sup>Department of Chemistry, Perm State University, 15 Bukireva St., Perm 614990, Russia; <sup>2</sup>Department of Technology for Organic Synthesis, Ural Federal University of the first President of Russia B. N. Yeltsin, 19 Mira St., Yekaterinburg, 620002, Russia.

E-mail: zhulan-nice91@yandex.ru

**Abstract.** During the past few decades dipolar cycloaddition reactions have become a powerful and effective tool for the synthesis of a wide range of heterocyclic compounds. In particular, the 1,3-dipolar cycloaddition reactions are successfully used in organic and medical chemistry to obtain a variety of five-membered heterocycles and the total synthesis of natural and biologically active products.<sup>1,2</sup> However, in sharp contrast, there is relatively little reported information on 1,4-dipolar cycloaddition reactions, despite the fact that the basic principles of such processes were proposed by R. Husgen almost half a century ago. In this regard, over the past two decades, 1,4-dipolar cycloaddition reactions are the subject of considerable research attention and, along with the Diels – Alder 4+2-cycloaddition, offer unique and alternative methods for producing six-membered heterocycles.<sup>3</sup>

For the first time we have discovered a simple and highly efficient method for the generation of cyclic enolate-iminium 1,4-dipoles.<sup>4</sup> *via* unique thermal decomposition of easily available dipyrazolodioxadiazocines **2**. By this approach various substituted spiro pyrazolo[5,1-*d*][1,3,5]dioxazines **3** have been synthesized by unusual cycloconversion of dipyrazolodioxadiazocines in the presence of ketones with overall high yields (up to 85% yield). Moreover, the developed method is 100% atom economical and proceeds under metal-catalyst- and solvent-free conditions.



**Scheme 1.** Cycloconversion of dipyrazolodioxadiazocines in the presence of ketones

#### References

1. Padwa, A., Pearson, W. H. (2002). *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*. Wiley: New York.
2. Hashimoto, T. Maruoka, K. (2015). Recent Advances of Catalytic Asymmetric 1,3-Dipolar Cycloadditions. *Chem. Rev.* Vol. 115, Iss. 11, pp. 5366–5412.
3. Nair, V., Deepthi, A., Ashok, D., Raveendran, A. E., Paul, R. R. (2014). 1,4-Dipolar cycloadditions and related reactions. *Tetrahedron*. Vol. 70, Iss. 19, pp. 3085–3105.
4. Zhulanov, V. E., Dmitriev, M. V., Maslivets, A. N., Rubin, M. (2016). New method for *in situ* generation of enolate-iminium 1,4-dipoles for [4+2] and [4+1] dipolar heterocycloaddition reactions. *RSC Adv.* Vol. 6, Iss. 93, pp. 90239–90247.

The work was supported by Russian Science Foundation, **project № 19-13- 00290**.